

Reply to "Comment on 'New ansatz for metric operator calculation in pseudo-Hermitian field theory' "

Abouzeid M. Shalaby*

Physics Department, Faculty of Science, Mansoura University, Egypt.

Abstract

In this report, we reply to a recent comment by Carl M. Bender, Gregorio Beninca and Hugh F. Jones on our work 'New ansatz for metric operator calculation in pseudo-Hermitian field theory (Phys. Rev. D. 79, 107702 (2009)). In fact, they figured out that there exist sign errors in our work which led to the conclusion of the invalidity of the ansatz introduced in our work. Here, we show that, the ansatz is valid in $d+1$ space-time dimensions, which by itself is a new and very important result. The importance of the work comes from the fact that it is the first time to have a metric operator for a quantum field theory which is local in the fields as well as valid in $3+1$ space-time dimensions. Moreover, it is composed of the operators in the Hamiltonian itself which makes the Feynmann diagram calculations for the physical amplitudes go the same way as in conventional theories.

PACS numbers: 12.90.+b, 11.30.Er, 12.60.Cn

Keywords: non-Hermitian models, $\mathcal{PT}$ -symmetric theories, pseudo-Hermitian theories, metric operator.

*Electronic address: amshalab@mans.edu.eg

In a recent comment, Carl M. Bender, Gregorio Beninca and Hugh F. Jones [1] found out some sign errors in our work in Ref.[2] which led them to conclude that the ansatz introduced in our work in Ref.[2] does not work in any space-time dimensions. In this report, we will show that the ansatz is valid not only in 1 + 1 space-time dimensions but also in all the space-time dimensions and represents the first calculations for a metric operator in the real world for a non-trivial non-Hermitian field theory.

The calculations of the metric operator are indispensable for any physical calculations in a pseudo-Hermitian theory. An important property of the metric operator is that it is not unique [3, 4, 5, 6, 7, 8]. This gives us the freedom to conjecture a simple form that can be employed in the calculations of the physical amplitudes in a practical way. In Ref. [9], a form of the C operator is obtained for the $i\phi^3$ field theory in 1 + 1 dimensions. This form is non-local in the fields and includes complicated parameters. In our work in Ref. [2], we introduced a new ansatz for the metric operator which is local in the fields and of fixed parameters as well. The idea of the ansatz is that the metric operator is conjectured to be a functional of all the operators in the Hamiltonian. To advocate this nice and interesting idea, in the following we present the final results of the first order Q operator for the $i\phi^3$ field theory. Note that, in the following, when two gradient operators appear in some formula this means that there exists an inner product between them *i.e* $\nabla\nabla \equiv \nabla \cdot \nabla$.

Now, consider the Hamiltonian model of the form;

$$H = \frac{1}{2} ((\nabla\phi)^2 + \pi^2 + m^2\phi^2) + ig\phi^3, \quad (1)$$

where ϕ is the field operator, π is the conjugate momentum and g is the coupling constant. The metric operator can be assumed to have the form $\eta = \exp(-Q)$, where Q is Hermitian. Note that, although the Hamiltonian in Eq.(1) is non-Hermitian in a Hilbert space with the Dirac sense inner product, it is Hermitian in a Hilbert space endowed by the inner product $\langle n|\eta|n\rangle$.

To obtain Q , we rename the terms in the Hamiltonian as;

$$H = H_0 + gH_I$$

$$H_0 = \frac{1}{2} \int d^d x ((\nabla\phi)^2 + \pi^2 + m^2\phi^2),$$

$$H_I = i \int d^d x \phi^3,$$

d is the spatial dimension. In using the relation $H^\dagger = \eta H \eta^{-1}$, we get

$$\begin{aligned} H^\dagger &= \exp(-Q) H \exp(Q) = H + [-Q, H] + \frac{[-Q, [-Q, H]]}{2!} \\ &\quad + \frac{[-Q, [-Q, [-Q, H]]]}{3!} + \dots, \end{aligned}$$

where

$$Q = Q_0 + gQ_1 + g^2Q_2 + g^3Q_3 + \dots$$

Up to first order in the coupling, we have the operator equation;

$$-2H_I = [-Q_1, H_0]. \quad (2)$$

In Ref.[2], we conjectured Q_1 to have the form;

$$\begin{aligned} Q_1 &= C_1 \int d^d z \pi^3(z) + \frac{C_2}{3} \int d^d z (\pi(z)\phi^2(z) + \phi(z)\pi(z)\phi(z) + \phi^2(z)\pi(z)) \\ &\quad + \frac{C_3}{3} \int d^d z (\pi(z)\nabla\phi(z)\nabla\phi(z) + \nabla\phi(z)\pi(z)\nabla\phi(z) + \nabla\phi(z)\nabla\phi(z)\pi(z)). \end{aligned} \quad (3)$$

To get the parameters C_1 , C_2 and C_3 , we substitute this form of Q_1 in Eq. (2).

Now, in commuting the first term in Q_1 with the second term in H_0 we get;

$$\begin{aligned} &C_1 \int d^d x \int d^d z \left[\pi^3(z), \frac{1}{2} \nabla_x \phi(x) \nabla_x \phi(x) \right] \\ &= \frac{3i}{2} C_1 \int d^d x \pi^2(x) \nabla_x^2 \phi(x) + \frac{3i}{2} C_1 \int d^d x \nabla_x^2 \phi(x) \pi^2(x), \end{aligned}$$

where it differs from the result in Ref.[2] by an overall negative sign.

For the first term in Q_1 with the third in H_0 , we have

$$\begin{aligned} &\frac{1}{2} m^2 C_1 \int d^d x \int d^d z [\pi^3(z), \phi(x) \phi(x)] \\ &= \frac{-3i}{2} m^2 C_1 \int d^d x \pi^2(x) \phi(x) + \frac{-3i}{2} m^2 C_1 \int d^d x \phi(x) \pi^2(x). \end{aligned}$$

Now, for the second term in Q_1 with the first in H_0 , we then have;

$$\begin{aligned} &\frac{C_2}{6} \int d^d x \int d^d z [\pi(z)\phi^2(z) + \phi(z)\pi(z)\phi(z) + \phi^2(z)\pi(z), \pi^2(x)] \\ &= iC_2 \int d^d x \phi(x) \pi^2(x) + iC_2 \int d^d x \pi^2(x) \phi(x) \end{aligned}$$

Also, for the second term in Q_1 with the second in H_0 , we have;

$$\begin{aligned} & \frac{C_2}{6} \int d^d x \int d^d z [\pi(z)\phi^2(z) + \phi(z)\pi(z)\phi(z) + \phi^2(z)\pi(z), \nabla_x \phi(x) \nabla_x \phi(x)] \\ &= iC_2 \int d^d x \phi^2(x) \nabla_x^2 \phi(x), \end{aligned} \quad (4)$$

where it differs from the result in Ref.[2] by an overall negative sign.

For the second term in Q_1 with the third in H_0 , we have

$$\begin{aligned} & \frac{C_2 m^2}{6} \int d^d x \int d^d z [\pi(z)\phi^2(z) + \phi(z)\pi(z)\phi(z) + \phi^2(z)\pi(z), \phi(x) \phi(x)] , \\ &= -iC_2 m^2 \int d^d x \phi^3(x). \end{aligned} \quad (5)$$

For the commutator of the third term in Q_1 with the first in H_0 ;

$$\begin{aligned} & \frac{C_3}{6} \int d^d x \int d^d z [\nabla_z \phi(z) \nabla_z \phi(z) \pi(z) + \nabla_z \phi(z) \pi(z) \nabla_z \phi(z) + \pi(z) \nabla_z \phi(z) \nabla_z \phi(z), \pi^2(x)] , \\ &= -iC_3 \int d^d x \pi(x) \pi(x) \nabla^2 \phi(x) - iC_3 \int d^d x \nabla^2 \phi(x) \pi(x) \pi(x). \end{aligned} \quad (6)$$

For the third of Q_1 with the second in H_0 we get;

$$\begin{aligned} & \frac{C_3}{6} \int d^d x \int d^d z \left[\begin{aligned} & \nabla_z \phi(z) \nabla_z \phi(z) \pi(z) + \nabla_z \phi(z) \pi(z) \nabla_z \phi(z) \\ & + \pi(z) \nabla_z \phi(z) \nabla_z \phi(z), \nabla_x \phi(x) \nabla_x \phi(x) \end{aligned} \right] \\ &= iC_3 \int d^d x \nabla^2 \phi(x) \nabla \phi(x) \nabla \phi(x). \end{aligned} \quad (7)$$

Also, for the third of Q_1 with the third in H_0 we get;

then

$$\begin{aligned} & \frac{m^2 C_3}{6} \int d^d x \int d^d z \left[\begin{aligned} & \nabla_z \phi(z) \nabla_z \phi(z) \pi(z) + \nabla_z \phi(z) \pi(z) \nabla_z \phi(z) \\ & + \pi(z) \nabla_z \phi(z) \nabla_z \phi(z), \phi(x) \phi(x) \end{aligned} \right] \\ &= \frac{im^2 C_3}{2} \int d^d x \nabla^2 \phi(x) \phi^2(x). \end{aligned} \quad (8)$$

For Eq.(2) to be satisfied, we have the following algebraic equations;

$$\begin{aligned}\frac{3i}{2}C_1 - iC_3 &= 0, \\ \frac{-3i}{2}m^2C_1 + iC_2 &= 0, \\ -iC_2m^2 &= 2i, \\ iC_2 + \frac{im^2C_3}{2} &= 0.\end{aligned}$$

Moreover, we have the integral $\int d^d x \nabla^2 \phi(x) \nabla \phi(x) \nabla \phi(x)$ which has to vanish as well. In fact, the above set of equations are inconsistent as they are four different equations in three unknowns. Nevertheless, one can verify the validity of the ansatz. To show this, we consider the fact that the quantized fields have to satisfy the Klein-Gordon equation because we have the relation;

$$\dot{\pi} = -i [\pi, H] = (\nabla^2 - m^2 - 3ig\phi) \phi, \quad (9)$$

which is nothing but the Klein-Gordon equation for the field ϕ . This non-linear equation can be solved exactly in the static frame with the solution given by [10];

$$\phi(x, y, z) = \frac{im^2}{2g} \text{sech}^2 \left(\frac{m(x + y + z)}{2\sqrt{3}} \right).$$

Since the field $\phi(x)$ transforms as $\phi(\Lambda x)$ under Lorentz group, one can get the exact solution at any frame of reference. With no loss of generality, let us assume the motion to be in the x -direction with a velocity v , then one has the solution;

$$\phi(x, y, z, t) = \frac{im^2}{2g} \text{sech}^2 \left(\frac{m(\gamma(x - vt) + y + z)}{2\sqrt{3}} \right),$$

where γ is the Lorentz factor. This solution has the following properties;

$$\begin{aligned}\nabla^2 \phi &= (3 - 2v^2) \frac{\partial^2 \phi}{\partial x^2}, \\ \frac{\partial \phi}{\partial y} &= \frac{\partial \phi}{\partial z} = \sqrt{1 - v^2} \frac{\partial \phi}{\partial x}, \\ \pi &= \frac{\partial \phi}{\partial t} = -v \frac{\partial \phi}{\partial x}.\end{aligned}$$

Accordingly we have;

$$\begin{aligned}\nabla^2 \phi(x) (\nabla \phi(x))^2 &= (v^2 - 2) (-3 + 2v^2) \left(\frac{\partial \psi}{\partial x} \right)^2 \left(\frac{\partial^2 \psi}{\partial x^2} \right), \\ \nabla^2 \phi(x) \pi^2(x) &= (3 - 2v^2) v^2 \left(\frac{\partial \psi}{\partial x} \right)^2 \left(\frac{\partial^2 \psi}{\partial x^2} \right),\end{aligned}$$

and since the integral $\int dx \frac{\partial^2 \phi(x)}{\partial x^2} \left(\frac{\partial \phi(x)}{\partial x} \right)^2$ is a vanishing surface term then the integrals $\int d^d x \nabla^2 \phi(x) \nabla \phi(x) \nabla \phi(x)$ and $\int d^d x \nabla^2 \phi(x) \pi^2(x)$ also vanish. Accordingly, for Eq.(2) to be satisfied we only have the following three algebraic equations for the parameters C_1 , C_2 and C_3 ;

$$\begin{aligned} \frac{-3i}{2} m^2 C_1 + i C_2 &= 0, \\ -i C_2 m^2 &= 2i, \\ i C_2 + \frac{i m^2 C_3}{2} &= 0. \end{aligned}$$

In other words, we have the parameters values as $C_1 = -\frac{4}{3m^4}$, $C_2 = -\frac{2}{m^2}$, $C_3 = \frac{4}{m^4}$, which is the same result reported in our work in Ref. [2] except that C_3 has reversed its sign.

To conclude, we have shown that the ansatz introduced in our work [2] is applicable in any space-time dimensions although of the sign errors mentioned in Ref.[1]. To show this, we relied on the properties of the exact form of the field which assured the validity of the ansatz. In fact, our ansatz is an important starting point for a concrete formulation of pseudo-Hermitian field theories since it is local in the fields and of simple shape. Moreover, the ansatz represents the first calculation for the metric operator of a non-trivial pseudo-Hermitian scalar field theory in the real world of $3 + 1$ space-time dimensions.

-
- [1] Carl M. Bender¹, Gregorio Beninca and H. F. Jones, Comment on “New ansatz for metric operator calculation in pseudo-Hermitian field theory” hep-th/09064293.
- [2] Abouzeid shalaby, Phys. Rev. D. 79, 107702 (2009)
- [3] C. M. Bender and S. P. Klevansky, Phys. Lett. A 373, 2670 (2009).
- [4] Ali Mostafazadeh, J. Phys. A: Math. Gen. 39 (2006) 10171–10188.
- [5] Mostafazadeh A 2002 Nucl. Phys. B 640 419.
- [6] Mostafazadeh A 2003 J. Math. Phys. 44 974.
- [7] Geyer H B, Scholtz F G and Snyman I 2004 Czech J. Phys. 54 1069.
- [8] Jones H F 2005 J. Phys. A: Math. Gen. 38 174.
- [9] C. M. Bender, D. C. Brody and H. F. Jones, Phys. Rev. Lett. 93, 251601 (2004); Phys. Rev. D70, 025001 (2004).
- [10] The details of how to obtain this exact solution will be presented in another work (in progress) but at the moment one can easily check that it really verify the Klein-Gordon equation.